

Executive Summary: Society of Critical Care Medicine Guidelines for the Treatment of Heat Stroke

KEYWORDS: cold water immersion; cooling methods; emergency medicine; heat stroke; heat-related illness; intensive care unit; medication prophylaxis; practice guideline

Heat-related illness is a leading cause of weather-related fatalities, and its prevalence is expected to increase (1). Heat stroke, the most severe form of heat-related illness, is defined as a core temperature greater than 40°C presenting with CNS abnormalities. Heat stroke can affect anyone, but certain populations are at higher risk including infants, elderly, athletes, domestically displaced, and outdoor workers (e.g., firefighters, construction workers, landscapers, military personnel, etc.) (2).

The increasing prevalence of heat stroke has emphasized the need for timely critical care and emergency services (3). Rapid cooling is an essential intervention to restore normal physiologic activity and minimize morbidity and mortality (4). Cold water immersion (CWI) is frequently cited as the preferred modality, but CWI may not be feasible in some settings or during mass casualty events, raising questions on the best alternative strategy (5). Additionally, once cooling has been achieved, critical care management of heat stroke is often centered around maintenance of organ support with minimal direct evidence to guide clinical decision-making. This executive summary provides an overview of the recommendations crafted by an international panel of multidisciplinary clinicians for the management of heat stroke. The seven Population, Intervention, Comparison, and Outcome (PICO) questions included in the guideline are presented in **Table 1**. For a detailed description of methodology, results, and rationale, the reader is referred to the complete guideline (6).

RECOMMENDATIONS

The panel issued a total of two strong recommendations, five good practice statements, and one “only-in-the-context of research” statement for these clinical practice guidelines (6). The infographic provides a brief summary of these recommendations (**Fig. 1**).

Recommendation 1

We recommend active cooling methods over passive cooling in patients with heat stroke (strong recommendation, very low certainty of evidence).

Remarks

CWI or ice water immersion will result in the fastest rate of temperature reduction and shortest time to target temperature (< 39°C).

Jeffrey F. Barletta, PharmD,
FCCM, FCCP¹

Tina L. Palmieri, MD, MCCM,
FACS, FABA²

Shari A. Toomey, MBA, RRT-NPS,
FCCM³

Fayez AlShamsi, MBBS⁴

Rebecca L. Stearns, PhD, ATC⁵

Asad E. Patanwala, PharmD,
MPH⁶

Nicole F. Siparsky, MD, FACS,
FCCM⁷

Neeraj Badjatia, MD, MSc,
MHCDs, FCCM⁸

Brian Schultz, MD, FAAP⁹

Crystal M. Breighner, MD, FACP¹⁰

Eric Bruno, MD, FAAEM¹¹

Christopher G. Harrod, MS¹²

Tanya Trevilian, MSN, RN, CCRN,
CPN, TCRN¹³

Leandro Braz de Carvalho, MD,
FCCM¹⁴

James Houser, MSN, RN, APRN,
FNP-C, NRP, CFRN, CMTE¹⁵

John M. Harahus, RN, NHDP-BC,
CHEC, CFRN, CEN¹⁶

Yang Liu, MD¹⁷

Ryan Swoboda¹⁸

Paulin Ruhato Banguti, MD¹⁹

Heatherlee Bailey, MD, MCCM,
FAAEM²⁰

Copyright © 2025 by the Society of Critical Care Medicine. All Rights Reserved.

DOI: 10.1097/CCM.0000000000006550

TABLE 1.**Population, Intervention, Comparison, and Outcome Questions and Summary of Recommendations**

Population, Intervention, Comparison, and Outcome Question	Recommendation
Population: Patients with heat stroke Intervention: Active cooling methods Comparison: Passive cooling methods Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, hospital survival, rate of temperature reduction, time to target temperature, achievement of target temperature within 30 min, organ dysfunction, adverse effects/complications, or hospital length of stay	We recommend active cooling methods over passive cooling in patients with heat stroke (strong recommendation; very low certainty of evidence)
Population: Patients with heat stroke Intervention: CWI (or other active method if not CWI) Comparison: Alternative method Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, hospital survival, rate of temperature reduction, time to target temperature, achievement of target temperature within 30 min, organ dysfunction, adverse effects/complications, or hospital length of stay	Clinicians should prioritize cooling methods that achieve the most rapid rate of cooling, which is ice water immersion (1–5°C) or CWI (9–12°C) (Good Practice Statement)
Population: Patients with classic heat stroke Intervention: CWI Comparison: Method other than CWI Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, hospital survival, rate of temperature reduction, time to target temperature, achievement of target temperature within 30 min, organ dysfunction, adverse effects/complications, or hospital length of stay	Clinicians may use similar cooling strategies for either classic or exertional heat stroke (Good Practice Statement)
Population: Patients with exertional heat stroke Intervention: CWI Comparison: Method other than CWI Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, hospital survival, rate of temperature reduction, time to target temperature, achievement of target temperature within 30 min, organ dysfunction, adverse effects/complications, or hospital length of stay	Clinicians may use similar cooling strategies for either classic or exertional heat stroke (Good Practice Statement)
Population: Patients with heat stroke Intervention: Reaching the target temperature within 30 min Comparison: Not reaching the target temperature within 30 min Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, survival, organ dysfunction, adverse effects/complications, or hospital length of stay	Clinicians should choose cooling methods that reach the target temperature within 30 min of recognition of heat stroke symptoms (Good Practice Statement)

(Continued)

TABLE 1. (Continued)**Population, Intervention, Comparison, and Outcome Questions and Summary of Recommendations**

Population, Intervention, Comparison, and Outcome Question	Recommendation
Population: Patients with heat stroke Intervention: Achieving a faster cooling rate ($\geq 0.155^{\circ}\text{C}/\text{min}$) Comparison: Not achieving a faster cooling rate ($< 0.155^{\circ}\text{C}/\text{min}$) Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, survival, organ dysfunction, adverse effects/complications, or hospital length of stay	Clinicians should prioritize cooling modalities that achieve a cooling rate $\geq 0.155^{\circ}\text{C}/\text{min}$ (Good Practice Statement)
Population: Patients with heat stroke Intervention: Medications that either directly or indirectly affect temperature control Comparison: No medications Outcome: Survival with good neurologic function long-term (i.e., after 6 mo), survival with good neurologic function at discharge, hospital survival, rate of temperature reduction, time to target temperature, achievement of target temperature within 30 min, organ dysfunction, adverse effects/complications, or hospital length of stay	We recommend against the use of dantrolene in patients with heat stroke (strong recommendation, very low certainty of evidence) The routine use of acetaminophen, non-steroidal anti-inflammatory drugs, and salicylates for temperature reduction should be avoided (Good Practice Statement) Prophylactic antibiotics or prophylactic antiseizure medications should only be used in the context of research (Only in the Context of Research Statement)

CWI = cold water immersion.

The outcome that was emphasized for this PICO was cooling rate because there were insufficient data to compare active vs. passive cooling methods on clinical outcomes such as survival, length of stay, organ dysfunction, and complications. A meta-analysis of randomized controlled trials (RCTs) (7–39) was performed that pooled the cooling rates for all active cooling methods and demonstrated a faster cooling rate with active vs. passive cooling in participants with experimental exertional hyperthermia (mean difference = $0.04^{\circ}\text{C}/\text{min}$ [95% CI, 0.03 – $0.05^{\circ}\text{C}/\text{min}$]). The overall certainty of evidence for the recommendation is very low, primarily due to indirectness (experimental exertional hyperthermia vs. actual heat stroke subjects), imprecision, and inconsistency.

Active cooling can be performed via numerous mechanisms, and there is substantial variability in the cooling rates achieved with each. For example, ice water and CWI have a cooling rate of approximately $0.178^{\circ}\text{C}/\text{min}$ and $0.154^{\circ}\text{C}/\text{min}$, respectively, compared with methods such as fan cooling ($0.057^{\circ}\text{C}/\text{min}$)

or evaporative cooling ($0.002^{\circ}\text{C}/\text{min}$) which are much slower. The CNS is extremely vulnerable to excessive heat and both the degree and duration of hyperthermia affect normal physiologic processes. In fact, cell death increases in an exponential manner as temperature exposure time increases (40). Data directly comparing cooling rates on clinical outcomes though are limited. One review combined case series data (not amenable to Grading of Recommendations, Assessment, Development, and Evaluation methodology) and reported higher mortality and a higher rate of complications when cooling rates of greater than $0.15^{\circ}\text{C}/\text{min}$ were not achieved (41). Similarly, two large case series reported zero fatalities when target temperatures were reached within 30 minutes (primarily with CWI) (42, 43). The panel decided active methods that achieve a cooling rate of at least $0.155^{\circ}\text{C}/\text{min}$ should be prioritized over those that do not achieve this rate. These include ice water immersion and CWI. CWI/ice water immersion may be associated with undesirable effects for patients,

2025 SCCM Guideline for the Treatment of Heat Stroke

Society of
Critical Care Medicine
The National Core Proficiency

- **Heat-related illness** is a *leading cause of weather-related fatalities* that are increasing in prevalence.
- **Heat stroke**, the most severe form of heat-related illness, is defined as a core temperature $>40^{\circ}\text{C}$ with central nervous system symptoms & can lead to significant morbidity & mortality.



Methods: An international team of multidisciplinary clinicians was formed to develop evidence-based guidelines, using GRADE methodology, on the management of heat stroke.

PICO:



1.) Cooling modalities



2.) Medications to reduce temperature

1. We recommend active cooling methods over passive cooling in patients with heat stroke.


[Strong recommendation; very low certainty of evidence]

- Prioritize cooling methods that achieve the most rapid rate of cooling which is ice- or cold-water immersion.
Good Practice Statement
- Use similar cooling strategies for either classic or exertional heat stroke.
Good Practice Statement
- Prioritize cooling modalities that achieve a cooling rate $\geq 0.155^{\circ}\text{C}/\text{min}$ and reach the target temperature within 30 min from recognition.
Good Practice Statements

2. We recommend against the use of dantrolene in patients with heat stroke.

[Strong recommendation; very low certainty of evidence]

- Avoid routine use of acetaminophen, NSAIDs & salicylates for temperature reduction.
Good Practice Statement
- Prophylactic antibiotics or prophylactic antiseizure medications should only be used in the context of research.
Only in the Context of Research Statement



- The figure above displays the weighted mean cooling rate ($^{\circ}\text{C}/\text{min}$) for each available cooling method, from fastest (left) to slowest (right).
- The table below displays each cooling method and the final core temperature achieved with that cooling method after 30 minutes for a given initial core temperature. Target final core temperatures ($<39^{\circ}\text{C}$) are colored green and near-target final core temperatures ($39-40^{\circ}\text{C}$) are colored yellow.

Cooling Method	Initial Core Temperature ($^{\circ}\text{C}$)						
	40.0	40.6	41.1	41.7	42.2	42.8	43.3
Final Core Temperature ($^{\circ}\text{C}$)							
Ice Water Immersion (1-5°C)	34.7	35.3	35.8	36.4	36.9	37.5	38.0
Colder Water Immersion (9-12°C)	35.4	36.0	36.5	37.1	37.6	38.2	38.7
Cold Water Immersion (14-17°C)	36.9	37.5	38.0	38.6	39.1	39.7	40.2
Tarp with Cold Water (approx. 25°C)	37.7	38.3	38.8	39.4	39.9	40.5	41.0
Cold Shower (approx. 21°C)	37.9	38.5	39.0	39.6	40.1	40.7	41.2
Ice Sheet Cooling	38.2	38.8	39.3	39.9	40.4	41.0	41.5
Cooling Vest	38.2	38.8	39.3	39.9	40.4	41.0	41.5
Hand Cooling Devices	38.3	38.9	39.4	40.0	40.5	41.1	41.6
Fan Cooling	38.3	38.9	39.4	40.0	40.5	41.1	41.6
Cold Water Immersion (14-17°C) of the Hands and/or Feet	38.4	39.0	39.5	40.1	40.6	41.2	41.7
Cold IV Saline	38.8	39.4	39.9	40.5	41.0	41.6	42.1
Commercial Ice Packs (neck, axilla, groin)	39.4	40.0	40.5	41.1	41.6	42.2	42.7
Reflective Blanket	39.4	40.0	40.5	41.1	41.6	42.2	42.7
Evaporative Cooling (Misting and Fanning)	39.9	40.5	41.0	41.6	42.1	42.7	43.2

Figure 1. Summary of guidelines recommendations for the management of heat stroke. To convert Celsius to Fahrenheit, multiply by 1.8 and add 32. GRADE = Grading of Recommendations, Assessment, Development, and Evaluation, NSAIDs = nonsteroidal anti-inflammatory drugs, SCCM = Society of Critical Care Medicine.

introduce hazards for hospital staff (e.g., slips and falls), complicate patient monitoring (e.g., cardiac monitor leads falling off), and interfere with ongoing resuscitation and procedures. Nevertheless, when considering the consequences of prolonged exposure to extreme body temperatures (i.e., long-term cognitive dysfunction and/or death), the panel determined that the desirable effects far outweighed the undesirable effects. CWI (or other methods of active cooling) may not be available or feasible in some settings such as austere environments, mass casualty events, or resource-limited regions. In these instances, clinicians are advised to select the method or combination of methods that will achieve the fastest rate of temperature reduction, aiming to achieve the target temperature within 30 minutes of recognition of heat stroke symptoms (see full online guideline). Careful planning and foresight are needed so required resources are readily available.

Recommendation 2

We recommend against the use of dantrolene in patients with heat stroke (strong recommendation, very low certainty of evidence).

Dantrolene is a treatment option for patients with malignant hyperthermia and its use in heat stroke has been considered due to overlap in pathophysiologic mechanisms. Three RCTs have compared dantrolene with placebo in patients with heat stroke (44–46). One is available in abstract form only. Meta-analyses have demonstrated no reduction in the risk of mortality (three trials; risk ratio [RR], 1.01 [95% CI, 0.11–9.31]), a mean difference in cooling time (two trials) of –10.5 minutes (95% CI, –28.5 to 7.6 min), a difference in hospital length of stay (one trial) of 1.85 days (95% CI, 0.99–2.70 d), no difference in the risk of adverse events (two trials; RR, 0.99 [95% CI, 0.46–2.15]), and a difference in incidence of recovery of consciousness as defined by a Glasgow Coma Score of greater than or equal to 13 points at less than or equal to 90 minutes of RR 2.5 (95% CI, 0.56–11.16). The overall certainty of evidence was very low due to risk of bias, imprecision, and inconsistency. When formulating the strength of the recommendation, the panel considered the quality of evidence, the wide CIs, including the fact that differences noted were not statistically significant, acquisition cost, and required

resources. Furthermore, some outcomes were only analyzed via one study.

Other medication therapies, such as acetaminophen, nonsteroidal anti-inflammatory drugs (NSAIDs), and salicylates, have no evidence supporting a desirable effect in the setting of heat stroke. Further, the panel recognized the risk for known adverse effects (e.g., hepatotoxicity, acute kidney injury, bleeding), which may be more prevalent in heat stroke patients. The routine use of acetaminophen, NSAIDs, and salicylates for temperature reduction should, therefore, be avoided. Due to the lack of any available evidence, antibiotic prophylaxis and seizure prophylaxis should only be used in the context of research.

RESEARCH AGENDA

Several challenges exist with conducting ethical clinical trials in patients with heat stroke. Given these challenges, most data arise from experiments conducted using healthy volunteers in controlled settings. Data utilizing endpoints like survival, survival with good neurologic function, organ failure, etc. are lacking. Further studies are needed that report outcomes in patients with heat stroke (vs. experimental hyperthermia) evaluating the relationship between cooling rate, time to temperature target, and clinical outcomes. Finally, research focused on treatment strategies in the critical care setting for comatose/symptomatic patients post-cooling is needed. A blueprint for future studies has been proposed (5).

- 1 Department of Pharmacy Practice, Midwestern University, College of Pharmacy, Glendale, AZ.
- 2 Davis and Shriners Children's Northern California, University of California, Sacramento, CA.
- 3 Carilion Rockbridge Community Hospital, Lexington, VA.
- 4 Department of Internal Medicine, College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates.
- 5 Korey Stringer Institute, University of Connecticut, Storrs, CT.
- 6 Faculty of Medicine and Health, Sydney School of Pharmacy, The University of Sydney, Sydney, NSW, Australia.
- 7 Rush University Medical Center, Chicago, IL.
- 8 Program in Trauma, Department of Neurology, University of Maryland School of Medicine, Baltimore, MD.

- 9 Brooke Army Medical Center, US Air Force, San Antonio, TX.
- 10 South Dayton Acute Care Consultants, Dayton, OH.
- 11 Department of Emergency Medicine, University of Tennessee Health Science Center, Murfreesboro, TN.
- 12 Society of Critical Care Medicine, Mount Prospect, IL.
- 13 Carilion Children's, Roanoke, VA.
- 14 Sociedade Mineira de Terapia Intensiva, Belo Horizonte, Brazil.
- 15 UPMC Center for Emergency Medicine of Western Pennsylvania, Pittsburgh, PA.
- 16 Geisinger Health System, Danville, PA.
- 17 Nanjing Gaochun People's Hospital, Nanjing, China.
- 18 Patient representative.
- 19 University of Rwanda and African Health Sciences University/King Faisal Hospital, Kigali, Rwanda.
- 20 Department of Emergency Medicine, Durham Veterans Affairs Medical Center, Durham, NC.

Dr. Barletta is a consultant for Wolters Kluwer and a current member of Council for the Society of Critical Care Medicine. Dr. Stearns is employed by the Korey Stringer Institute, member of the USA Football Medical Advisory Panel, and received royalties as book editor for two books that contain material on heat illness. Dr. Siparsky received funding from UpToDate. Dr. Schultz disclosed government work; he is a U.S. Government employee. Dr. Bruno received funding from Hemcon. Dr. Bailey received funding from Radiometer and Baxter; she disclosed that she is a past President of the Society of Critical Care Medicine. The remaining authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: jbarle@midwestern.edu

REFERENCES

1. National Oceanic and Atmospheric Administration/National Weather Service: Weather Related Fatality and Injury Statistics. 2023. Available at: [weather.gov/hazstat](https://www.weather.gov/hazstat). Accessed August 15, 2024
2. Douma MJ, Aves T, Allan KS, et al; First Aid Task Force of the International Liaison Committee on Resuscitation: First aid cooling techniques for heat stroke and exertional hyperthermia: A systematic review and meta-analysis. *Resuscitation* 2020; 148:173–190
3. National EMS Information System: Heat-Related EMS Activation Surveillance Dashboard. 2024. Available at: <https://nemsis.org/heat-related-ems-activation-surveillance-dashboard/>. Accessed August 15, 2024
4. Bouchama A, Abuyassin B, Lehe C, et al: Classic and exertional heatstroke. *Nat Rev Dis Primers* 2022; 8:8
5. Barletta JF, Palmieri TL, Toomey SA, et al: Management of heat-related illness and injury in the ICU: A concise definitive review. *Crit Care Med* 2024; 52:362–375
6. Barletta JF, Palmieri TL, Toomey SA, et al: Society of Critical Care Medicine Guideline for the Treatment of Heat Stroke. *Crit Care Med* 2025; 53:e490–e500
7. Clements JM, Casa DJ, Knight J, et al: Ice-water immersion and cold-water immersion provide similar cooling rates in runners with exercise-induced hyperthermia. *J Athl Train* 2002; 37:146–150
8. DeMartini JK, Ranalli GF, Casa DJ, et al: Comparison of body cooling methods on physiological and perceptual measures of mildly hyperthermic athletes. *J Strength Cond Res* 2011; 25:2065–2074
9. Peiffer JJ, Abbiss CR, Watson G, et al: Effect of cold-water immersion duration on body temperature and muscle function. *J Sports Sci* 2009; 27:987–993
10. Peiffer JJ, Abbiss CR, Watson G, et al: Effect of a 5-min cold-water immersion recovery on exercise performance in the heat. *Br J Sports Med* 2010; 44:461–465
11. Taylor NA, Caldwell JN, Van den Heuvel AM, et al: To cool, but not too cool: That is the question—immersion cooling for hyperthermia. *Med Sci Sports Exerc* 2008; 40:1962–1969
12. Walker A, Driller M, Brearley M, et al: Cold-water immersion and iced-slush ingestion are effective at cooling firefighters following a simulated search and rescue task in a hot environment. *Appl Physiol Nutr Metab* 2014; 39:1159–1166
13. Weiner JS, Khogali M: A physiological body-cooling unit for treatment of heat stroke. *Lancet* 1980; 1:507–509
14. Barwood MJ, Davey S, House JR, et al: Post-exercise cooling techniques in hot, humid conditions. *Eur J Appl Physiol* 2009; 107:385–396
15. Carter JM, Rayson MP, Wilkinson DM, et al: Strategies to combat heat strain during and after firefighting. *J Therm Biol* 2007; 32:109–116
16. Clapp AJ, Bishop PA, Muir I, et al: Rapid cooling techniques in joggers experiencing heat strain. *J Sci Med Sport* 2001; 4:160–167
17. Selkirk GA, McLellan TM, Wong J: Active versus passive cooling during work in warm environments while wearing firefighting protective clothing. *J Occup Environ Hyg* 2004; 1:521–531
18. Zhang Y, Nepocatyck S, Katika CP, et al: Effects of half time cooling on thermoregulatory responses and soccer-specific performance tests. *Monten J Sports Sci Med* 2014; 3:17–22
19. Halson SL, Quod MJ, Martin DT, et al: Physiological responses to cold water immersion following cycling in the heat. *Int J Sports Physiol Perform* 2008; 3:331–346
20. Hosokawa Y, Adams WM, Belval LN, et al: Tarp-assisted cooling as a method of whole-body cooling in hyperthermic individuals. *Ann Emerg Med* 2017; 69:347–352
21. Butts CL, McDermott BP, Buening BJ, et al: Physiologic and perceptual responses to cold-shower cooling after exercise-induced hyperthermia. *J Athl Train* 2016; 51:252–257
22. Flouris AD, Wright-Beatty HE, Friesen BJ, et al: Treatment of exertional heat stress developed during low or moderate physical work. *Eur J Appl Physiol* 2014; 114:2551–2560
23. Gagnon D, Lemire BB, Casa DJ, et al: Cold-water immersion and the treatment of hyperthermia: Using 38.6 degrees C as a safe rectal temperature cooling limit. *J Athl Train* 2010; 45:439–444
24. Luhring KE, Butts CL, Smith CR, et al: Cooling effectiveness of a modified cold-water immersion method after exercise-induced hyperthermia. *J Athl Train* 2016; 51:946–951
25. Butts CL, Spisla DL, Adams JD, et al: Effectiveness of ice-sheet cooling following exertional hyperthermia. *Mil Med* 2017; 182:e1951–e1957

26. Caldwell AR, Saillant MM, Pitsas D, et al: The effectiveness of a standardized ice-sheet cooling method following exertional hyperthermia. *Mil Med* 2022; 187:e1017–e1023
27. Pryor RR, Haboian K, Fitts T, et al: Tarp-assisted cooling for exertional heat stroke treatment in wildland firefighting. *Wilderness Environ Med* 2023; 34:490–497
28. Hosokawa Y, Belval LN, Adams WM, et al: Chemically activated cooling vest's effect on cooling rate following exercise-induced hyperthermia: A randomized counter-balanced crossover study. *Medicina (Kaunas)* 2020; 56:539
29. Lopez RM, Cleary MA, Jones LC, et al: Thermoregulatory influence of a cooling vest on hyperthermic athletes. *J Athl Train* 2008; 43:55–61
30. Smith CR, Butts CL, Adams JD, et al: Effect of a cooling kit on physiology and performance following exercise in the heat. *J Sport Rehabil* 2018; 27:413–418
31. Kielblock AJ, Van Rensburg JP, Franz RM: Body cooling as a method for reducing hyperthermia. An evaluation of techniques. *S Afr Med J* 1986; 69:378–380
32. Sefton JM, McAdam JS, Pascoe DD, et al: Evaluation of 2 heat-mitigation methods in army trainees. *J Athl Train* 2016; 51:936–945
33. Lissoway JB, Lipman GS, Grahn DA, et al: Novel application of chemical cold packs for treatment of exercise-induced hyperthermia: A randomized controlled trial. *Wilderness Environ Med* 2015; 26:173–179
34. Reynolds KA, Evanich JJ, Eberman LE: Reflective blankets do not effect cooling rates after running in hot, humid conditions. *Int J Exerc Sci* 2015; 8:97–103
35. Adams WM, Hosokawa Y, Adams EL, et al: Reduction in body temperature using hand cooling versus passive rest after exercise in the heat. *J Sci Med Sport* 2016; 19:936–940
36. Maroni T, Dawson B, Barnett K, et al: Effectiveness of hand cooling and a cooling jacket on post-exercise cooling rates in hyperthermic athletes. *Eur J Sport Sci* 2018; 18:441–449
37. Zhang Y, Bishop PA, Casaru C, et al: A new hand-cooling device to enhance firefighter heat strain recovery. *J Occup Environ Hyg* 2009; 6:283–288
38. McDermott BP, Atkins WC: Whole-body cooling effectiveness of cold intravenous saline following exercise hyperthermia: A randomized trial. *Am J Emerg Med* 2023; 72:188–192
39. Caldwell JN, van den Heuvel AMJ, Kerry P, et al: A vascular mechanism to explain thermally mediated variations in deep-body cooling rates during the immersion of profoundly hyperthermic individuals. *Exp Physiol* 2018; 103:512–522
40. Walter EJ, Carraretto M: The neurological and cognitive consequences of hyperthermia. *Crit Care* 2016; 20:199
41. Filep EM, Murata Y, Endres BD, et al: Exertional heat stroke, modality cooling rate, and survival outcomes: A systematic review. *Medicina (Kaunas)* 2020; 56:589
42. Stearns RL, Hosokawa Y, Belval LN, et al: Exertional heat stroke survival at the Falmouth Road Race: 180 New cases with expanded analysis. *J Athl Train* 2024; 59:304–309
43. Demartini JK, Casa DJ, Stearns R, et al: Effectiveness of cold water immersion in the treatment of exertional heat stroke at the Falmouth Road Race. *Med Sci Sports Exerc* 2015; 47:240–245
44. Bouchama A, Cafege A, Devol EB, et al: Ineffectiveness of dantrolene sodium in the treatment of heatstroke. *Crit Care Med* 1991; 19:176–180
45. Channa AB, Seraj MA, Saddique AA, et al: Is dantrolene effective in heat stroke patients? *Crit Care Med* 1990; 18:290–292
46. Hepner A, Greenberg M: Safety and efficacy of dantrolene sodium (250 mg/5 mL) in patients with exertional heat stroke. *Ann Emerg Med* 2017; 70:S39